arXiv:1110.2889v1 [math-ph] 13 Oct 2011

Multitime Rayleigh Solitons

Laura Gabriela Matei, Constantin Udriste

University POLITEHNICA of Bucharest, Faculty of Applied Sciences,
Department of Mathematics-Informatics I, Splaiul Independentei 313,
Bucharest 060042, Romania, E-mails: mateiglaura@yahoo.com;
udriste@mathem.pub.ro; anet.udri@yahoo.com

Abstract

Multitime evolution PDEs for Rayleigh waves are considered, using geometrical ingredients capable to build an ultra-parabolic-hyperbolic differential operator. Their soliton solutions are found based on appropriate hypotheses and Bernoulli ODEs. These multitime solitons develop complex behavior of deformation phenomena.

Section 1 presents the single-time Rayleigh wave equations. Section 2 analyzes the geometric characteristics (fundamental tensor, linear connection, vector fields, tensor fields) associated to the Rayleigh PDEs, showing the existence of an infinity of geometrical structures such that the multitime PDEs are prolongations of single-time PDE. Section 3 builds the ultra-parabolic-hyperbolic differential operator defining the multitime Rayleigh wave equations. Section 4 gives the technique capable to produce the multitime Rayleigh solitons. Sections 5 and 6 praise explicit formulas for the multitime Rayleigh solitons.

Mathematics Subject Classification 2010: 37K40, 35C08, 35Q51.
Key words: multitime solitons, single-time Rayleigh PDE, multitime Rayleigh PDE, Bernoulli ODE.

1 Single-time Rayleigh PDEs

In the physical and mathematical literature [1]-[4], [7]-[9], we find the *Rayleigh wave equation*

$$u_{tt} - u_{xx} = \epsilon(u_t - u_t^3) \tag{R}$$

related to *Rayleigh wave equation of Van der Pol type*

$$u_{tt} - u_{xx} = \epsilon(1 - u^2)u_t. \tag{RVP}$$

Each of these has been used to model physical phenomena. Now, the PDE (R) serves as a model for the large amplitude vibrations of wind-blown, ice-laden power transmission lines, in time that, the PDE (RVP) describes plane electromagnetic waves propagating between two parallel planes in a region where the conductivity varies quadratically with the electric field.

Just as their counterparts from ordinary differential equations, the PDEs (R) and (VP) can be transformed one to another. Their solutions can be obtained by simple operations performed on the solution of a certain first order, nonlinear wave equation.

An initial-boundary value problem for Rayleigh nonlinear wave equation can be considered to be a simple model to describe the galloping oscillations of overhead power transmission lines in a windfield. One end of the transmission line is assumed to be fixed, whereas the other end of the line is assumed to be attached to a dashpot system.

Rayleigh surface waves are of particular importance in seismology, acoustic, geophysics and electronics applications.

Remark Some papers [6] describe as Rayleigh wave a type of seismic surface wave that moves with a rolling motion that consists of a combination of particle motion perpendicular and parallel to the main direction of wave propagation. The amplitude of this motion decreases with depth. Like primary waves, Rayleigh waves are alternatingly compressional and extensional (they cause changes in the volume of the rocks they pass through).

2 Multitime PDE systems

The passing from systems of PDEs with a single-time variable t to related PDE systems with $m \geq 2$ evolution variables $t = (t^\alpha)$, $\alpha = 1, ..., m$, is substantially complicated due to necessity of praising some reasons and some techniques of such change. The most natural way of changing is to use geometrical ingredients (derivation, trace etc) that extend the initial system. The theory and a systematic procedure for the construction of such new systems is presented here in the context of Rayleigh nonlinear waves.

This paper provides new results regarding the multitime solitons in two and more temporal dimensions that can be of interest in physics. We overpass the complexity and, furthermore, the difficulty of performing hand computations for Rayleigh PDE systems involving many temporal variables by using the symbolic software package in MAPLE. Our source of inspiration for introducing and studying multitime soliton PDEs is the paper [5].

Also the papers [10]-[17] contains a lot of ideas in this direction.

3 Multitime geometrical prolongations of Rayleigh PDEs

Let us introduce and study some *multitime geometrical prolongations of the Rayleigh PDEs*, using related connection, fundamental tensor field, vector fields, tensor fields which leave on the jet bundle as ingredients in the Differential Geometry of the manifold R^m.

Suppose the *multitime* $t = (t^1, ..., t^m) \in \mathbb{R}^m$ is a parameter of evolution. We endow the manifold (jet bundle of order one) $J^1(\mathbb{R}^m \times \mathbb{R}, \mathbb{R} \times \mathbb{R}^m)$ with a *distinguished symmetric linear connection* $\Gamma^\gamma_{\alpha\beta} = \Gamma^\gamma_{\alpha\beta}\left(t, x, u, \frac{\partial u}{\partial t}\right)$, and with a *distinguished fundamental symmetric contravariant tensor field* $h = \left(h^{\alpha\beta}(t, x, u, \frac{\partial u}{\partial t})\right)$ of constant signature (r, z, s), $r + z + s = m$. Using a C^2 function $u : \mathbb{R}^m \times \mathbb{R} \to \mathbb{R}$, we build the *Hessian operator*

$$(Hess_\Gamma u)_{\alpha\beta} = \frac{\partial^2 u}{\partial t^\alpha \partial t^\beta} - \Gamma^\gamma_{\alpha\beta}\frac{\partial u}{\partial t^\gamma}, \quad \alpha, \beta, \gamma \in \{1, ..., m\}$$

and its trace, called *ultra-parabolic-hyperbolic operator*,

$$\Box_{\Gamma,h} u = h^{\alpha\beta}(Hess_\Gamma u)_{\alpha\beta}.$$

We define a *multitime PDE* as

$$\Box_{\Gamma,h} u - \frac{\partial^2 u}{\partial x^2} = 0, \tag{1}$$

where $t = (t^1, ..., t^m) \in R^m$ and $x \in R$.

- Let

$$C^\gamma(t, x, \eta, \xi),\ \gamma = 1, ..., m$$

be a *distinguished vector field* and

$$B^{\alpha\beta\gamma}(t, x, \eta, \xi),\ \alpha, \beta, \gamma \in \{1, ..., m\}$$

be a *distinguished tensor field.* If we adopt the hypothesis

$$h^{\alpha\beta}(t, x, \eta, \xi)\Gamma^\gamma_{\alpha\beta}(t, x, \eta, \xi)\xi_\gamma = C^\gamma(t, x, \eta, \xi)\xi_\gamma - B^{\alpha\beta\gamma}(t, x, \eta, \xi)\xi_\alpha\xi_\beta\xi_\gamma, \tag{2}$$

then we obtain the *multitime Rayleigh PDE*

$$h^{\alpha\beta}\frac{\partial^2 u}{\partial t^\alpha \partial t^\beta} - C^\gamma \frac{\partial u}{\partial t^\gamma} + B^{\alpha\beta\gamma}\frac{\partial u}{\partial t^\alpha}\frac{\partial u}{\partial t^\beta}\frac{\partial u}{\partial t^\gamma} - \frac{\partial^2 u}{\partial x^2} = 0. \tag{3}$$

- If $C^\gamma(t,x,\eta,\xi)$, $\gamma = 1,...,m$ and $D^\gamma(t,x,\eta,\xi)$, $\gamma = 1,...,m$ are *distinguished vector fields* and the constraint relation is

$$h^{\alpha\beta}(t,x,\eta,\xi)\Gamma^\gamma_{\alpha\beta}(t,x,\eta,\xi)\xi_\gamma = C^\gamma(t,x,\eta,\xi)\xi_\gamma - D^\gamma(t,x,\eta,\xi)\eta^2\xi_\gamma, \tag{2'}$$

then we get a *multitime Rayleigh wave equation of Van der Pol type*

$$h^{\alpha\beta}\frac{\partial^2 u}{\partial t^\alpha \partial t^\beta} - C^\gamma \frac{\partial u}{\partial t^\gamma} + u^2 D^\gamma \frac{\partial u}{\partial t^\gamma} - \frac{\partial^2 u}{\partial x^2} = 0. \tag{3'}$$

Theorem

(1) There exists an infinity of geometrical structures $\Gamma^\gamma_{\alpha\beta}$*,* $h^{\alpha\beta}$*,* C^γ*,* $B^{\alpha\beta\gamma}$ *on* R^m *such that a solution of the Rayleigh PDE (R) is also a solution of the multitime Rayleigh PDE (3).*

(2) There exists an infinity of geometrical structures $\Gamma^\gamma_{\alpha\beta}$*,* $h^{\alpha\beta}$*,* C^γ*,* D^γ *on* R^m *such that a solution of the Rayleigh PDE (RVP) is also a solution of the multitime Rayleigh PDE of Van der Pol type* $(3')$*.*

Proof Let $t^1 = t$ and $u = u(t^1, x)$.

(1) Suppose $u = u(t^1,x)$ is a solution of single-time Rayleigh PDE (R). The function $v(t^1,...,t^m,x) = u(t^1,x)$ is a solution of the multitime Rayleigh PDE (3) if the family of geometrical structures $\Gamma^\gamma_{\alpha\beta}$, $h^{\alpha\beta}$, C^γ, $B^{\alpha\beta\gamma}$ is fixed by

$$h^{\alpha\beta}\Gamma^1_{\alpha\beta}\xi_1 = C^1\xi_1 - B^{111}\xi_1\xi_1\xi_1.$$

It is obvious that we have an infinity of geometrical structures that satisfy this algebraic equation.

(2) Suppose $u = u(t^1,x)$ is a solution of single-time Rayleigh PDE (RVP). The function $v(t^1,...,t^m,x) = u(t^1,x)$ is a solution of the multitime Rayleigh PDE $(3')$ if the family of geometrical structures $\Gamma^\gamma_{\alpha\beta}$, $h^{\alpha\beta}$, C^γ, D^γ is fixed by

$$h^{\alpha\beta}\Gamma^1_{\alpha\beta}\xi_1 = C^1\xi_1 - D^1u^2\xi_1.$$

It is obvious that we have an infinity of geometrical structures that satisfy this algebraic equation.

The foregoing Theorem justifies the term *multitime geometrical prolongations of the Rayleigh PDEs.*

4 Multitime Rayleigh solitons

The first aim of this Section is to find some multitime solitons solutions for this multitime PDEs. In spite of the mathematical beauty, the distance between theoretical multitime models and real situations where they apply is far from our understanding.

Let $\phi : I \subset \mathbb{R} \to \mathbb{R}$ be a function of class C^4. We seek for solutions of the PDEs (3) and (3′) in the form of *multitime solitons*

$$u(x,t) = \phi(x - \lambda_\alpha t^\alpha) = \phi(z), \tag{4}$$

where λ_α, $\alpha = 1, ..., m$, is a constant vector and $z = x - \lambda_\alpha t^\alpha$. Then, the partial derivatives of the unknown function $u(x,t)$ are

$$\frac{\partial^2 u}{\partial x^2} = \phi''(z), \ \frac{\partial u}{\partial t^\alpha} = \phi'(z)(-\lambda_\alpha), \ \frac{\partial^2 u}{\partial t^\alpha \partial t^\beta} = \phi''(z)\lambda_\alpha\lambda_\beta. \tag{5}$$

Substituting these derivatives in the PDEs (3) and (3′), we obtain the second order ODEs,

$$[h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1]\,\phi''(z) - B^{\alpha\beta\gamma}\lambda_\alpha\lambda_\beta\lambda_\gamma\,\phi'(z)^3 + C^\gamma\lambda_\gamma\,\phi'(z) = 0 \tag{6}$$

and respectively

$$[h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1]\,\phi''(z) - D^\gamma\lambda_\gamma\,\phi^2(z)\phi'(z) + C^\gamma\lambda_\gamma\,\phi'(z) = 0. \tag{6′}$$

Summarizing, we have

Theorem *If $\phi(z)$ is a solution of second order ODE* (6) *or* (6′) *, then $u(x,t) = \phi(x - \lambda_\alpha t^\alpha)$ is a multitime soliton solution of the multitime Rayleigh PDEs (3) or (3′) respectively.*

In order to find some multitime Rayleigh solitons, we make some particular choices of the elements that appear in the construction of the two multitime PDEs.

First, for multitime Rayleigh solitons, we consider the ODE (6) and we relate the metric tensor $h^{\alpha\beta}(t,x,\eta,\xi)$, the tensor field $B^{\alpha\beta\gamma}(t,x,\eta,\xi)$, the vector field $C^\gamma(t,x,\eta,\xi)$, $\gamma = 1, ..., m$ and the constant vector λ_α by the conditions

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = a(x - \lambda_\alpha t^\alpha) = a(z) \neq 0, \\ B^{\alpha\beta\gamma}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta\lambda_\gamma = b(x - \lambda_\alpha t^\alpha) = b(z), \\ C^\gamma(t,x,\eta,\xi)\lambda_\gamma = c(x - \lambda_\alpha t^\alpha) = c(z). \end{cases}$$

With these conditions, the ODE (6) becomes

$$a(z)\phi'' - b(z)(\phi')^3 + c(z)\phi' = 0. \tag{7}$$

Second, for multitime Rayleigh solitons, we take de ODE (6′) and we relate the metric tensor $h^{\alpha\beta}(t,x,\eta,\xi)$, the vector fields $C^\gamma(t,x,\eta,\xi)$ and $D^\gamma(t,x,\eta,\xi)$, $\gamma = 1,...,m$ and the constant vector λ_α by the conditions

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = a(x-\lambda_\alpha t^\alpha) = a(z) \neq 0, \\ D^\gamma(t,x,\eta,\xi)\lambda_\gamma = b(x-\lambda_\alpha t^\alpha) = d(z), \\ C^\gamma(t,x,\eta,\xi)\lambda_\gamma = c(x-\lambda_\alpha t^\alpha) = c(z). \end{cases}$$

With this choice, the ODE (6′) becomes

$$a(z)\phi'' - d(z)\phi^2\phi' + c(z)\phi' = 0. \tag{7′}$$

We are looking for some solutions of the ODE (7) and (7′), with a view to finding multitime Rayleigh solitons.

5 Families of multitime Rayleigh solitons

5.1 Case of coefficients depending on z

Denoting $\phi' \overset{\text{not}}{=} \psi$, the second order ODE (7) becomes a first order ODE,

$$a(z)\,\psi' - b(z)\,\psi^3 + c(z)\,\psi = 0, \tag{8}$$

called *Bernoulli ODE*. The general form of this ODE is called *Abel ODE* of the *first kind*, and it arose in the context of the studies of Niels Henrik Abel on the theory of *elliptic functions*, and represents a natural generalization of the Riccati equation.

Since $a(z) \neq 0$, we write the equation in the form

$$\psi' = -\frac{c(z)}{a(z)}\,\psi + \frac{b(z)}{a(z)}\,\psi^3. \tag{9}$$

By a change of the unknown function, $\xi = \psi^{-2}$, the Bernoulli ODE becomes a linear ODE,

$$\xi' - 2\,\frac{c(z)}{a(z)}\,\xi = -2\,\frac{b(z)}{a(z)}, \tag{10}$$

with the solutions

$$\xi(z) = e^{2\int \frac{c(z)}{a(z)}dz} \left[K - 2\int \frac{b(z)}{a(z)}\, e^{-2\int \frac{c(z)}{a(z)}dz} dz \right], \; K \in \mathbb{R}. \tag{11}$$

Since $\psi = \xi^{-\frac{1}{2}}$, we obtain

$$\psi(z) = e^{-\int \frac{c(z)}{a(z)}dz} \left[K - 2\int \frac{b(z)}{a(z)}\, e^{-2\int \frac{c(z)}{a(z)}dz} dz \right]^{-\frac{1}{2}}, \; K \in \mathbb{R},$$

that is

$$\phi'(z) = \frac{e^{-\int \frac{c(z)}{a(z)}dz}}{\sqrt{K - 2\int \frac{b(z)}{a(z)}\, e^{-2\int \frac{c(z)}{a(z)}dz} dz}}, \; K \in \mathbb{R}. \tag{12}$$

Therefore, we have found solutions of the multitime PDE Rayleigh (3):
Theorem *If we fix the above elements by the conditions*

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = a(x - \lambda_\alpha t^\alpha) \neq 0, \\ B^{\alpha\beta\gamma}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta\lambda_\gamma = b(x - \lambda_\alpha t^\alpha), \\ C^\gamma(t,x,\eta,\xi)\lambda_\gamma = c(x - \lambda_\alpha t^\alpha), \end{cases}$$

then the function

$$u(x,t) = \phi(x - \lambda_\alpha t^\alpha)$$

represents a multitime soliton-solution for the multitime PDE Rayleigh (3) for every ϕ given by

$$\phi(z) = \int \frac{e^{-\int \frac{c(z)}{a(z)}dz}}{\sqrt{K - 2\int \frac{b(z)}{a(z)}\, e^{-2\int \frac{c(z)}{a(z)}dz} dz}}\, dz, \; K \in \mathbb{R}. \tag{13}$$

5.2 Case of constant coefficients

If we fix the elements h, B, C, λ by the conditions

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = \text{constant} = a \neq 0, \\ B^{\alpha\beta\gamma}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta\lambda_\gamma = \text{constant} = b, \\ C^{\gamma}(t,x,\eta,\xi)\lambda_\gamma = \text{constant} = c, \end{cases}$$

then the ODE (7) takes the form

$$a\phi'' - b(\phi')^3 + c\phi' = 0. \tag{14}$$

Denoting $\phi' \overset{\text{not}}{=} \psi$, the second order ODE with constant coefficients (14) becomes a first order ODE, called *Bernoulli ODE*

$$a\,\psi' - b\,\psi^3 + c\,\psi = 0. \tag{15}$$

The ODE (15) has separable variables, i.e., it can be written

$$a\,\frac{d\psi}{b\psi^3 - c\psi} = dz \;\;\Leftrightarrow\;\; \frac{a}{b}\int \frac{d\psi}{\psi\left(\psi^2 - \dfrac{c}{b}\right)} = \int dz$$

and we get the next sequence of equivalences

$$\frac{a}{c}\left(-\int \frac{d\psi}{\psi} + \int \frac{\psi d\psi}{\psi^2 - \dfrac{c}{b}}\right) = \int dz$$

$$\Leftrightarrow \ln\left(\frac{\sqrt{\left|\psi^2 - \dfrac{c}{b}\right|}}{|\psi|}\right) = \frac{c}{a}z + l, \;\; l \in \mathbb{R}$$

$$\Leftrightarrow \psi^2 - \frac{c}{b} = p\,\psi^2 e^{\frac{2c}{a}z}, \;\; p \in \mathbb{R}^*$$

$$\Leftrightarrow \psi(z) = \pm\sqrt{\frac{c}{b(1 - p\,e^{\frac{2c}{a}z})}}, \;\; p \in \mathbb{R}^*, \;\; \text{with} \;\; \frac{c}{b(1 - p\,e^{\frac{2c}{a}z})} \geq 0,$$

that is

$$\phi(z) = \pm\int\sqrt{\frac{c}{b(1 - p\,e^{\frac{2c}{a}z})}}\,dz, \;\; p \in \mathbb{R}^*, \;\; \text{with} \;\; \frac{c}{b(1 - p\,e^{\frac{2c}{a}z})} \geq 0. \tag{16}$$

To calculate the primitive from the right hand side, we amplify the fraction by $e^{\frac{2c}{a}z}$ and then make the change of variables $e^{\frac{c}{a}z} = t$. We obtain the integral

$$I = \pm \frac{a}{c} \int \frac{1}{t} \sqrt{\frac{c}{b(1-pt^2)}}\, dt$$

and a new change of variables, $\frac{1}{t} = s$, gives a new integral,

$$J = \pm \frac{a}{c} \int \sqrt{\frac{c}{b(s^2-p)}}\, ds.$$

There are two separate possibilities:

a) If we take $\frac{c}{b} > 0$, then

$$J = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \int \frac{1}{\sqrt{s^2-p}}\, ds.$$

- For $p > 0$, the integral is

$$J = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \operatorname{ch}^{-1} \left| \frac{s}{\sqrt{p}} \right| + r, \quad r \in \mathbb{R},$$

that is

$$\phi(z) = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \operatorname{ch}^{-1} \left| \frac{1}{e^{\frac{c}{a}z}\sqrt{p}} \right| + r, \quad r \in \mathbb{R}, \quad p \in \mathbb{R}_+^*$$

and we can write

$$\phi(z) = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \operatorname{ch}^{-1} \left(K e^{-\frac{c}{a}z} \right) + r, \quad r \in \mathbb{R}, \quad K \in \mathbb{R}_+^*. \tag{17}$$

- For $p < 0$, we have

$$J = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \int \sqrt{\frac{1}{s^2+p'}}\, ds, \quad p' = -p, \quad p' > 0.$$

It follows

$$J = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \operatorname{sh}^{-1} \left| \frac{s}{\sqrt{p'}} \right| + r, \quad r \in \mathbb{R},$$

that is

$$\phi(z) = \pm \frac{a}{c} \sqrt{\frac{c}{b}} \operatorname{sh}^{-1} \left| \frac{1}{e^{\frac{c}{a}z}\sqrt{p'}} \right| + r, \quad r \in \mathbb{R}, \quad p' \in \mathbb{R}_+^*$$

and then

$$\phi(z) = \pm \frac{a}{c}\sqrt{\frac{c}{b}}\ \mathrm{sh}^{-1}\left(Ke^{-\dfrac{c}{a}z}\right) + r, \quad r \in \mathrm{I\!R}, \quad K \in \mathrm{I\!R}^*_+. \tag{18}$$

b) If we suppose $\frac{c}{b} < 0$, then, via $\frac{c}{b(1 - p\, e^{\frac{2c}{a}z})} \geq 0$, it follows $p > 0$ and the integral J becomes

$$J = \pm \frac{a}{c}\sqrt{\frac{-c}{b}} \int \frac{1}{\sqrt{p - s^2}}\, ds, \quad p > 0$$

$$\Leftrightarrow J = \pm \frac{a}{c}\sqrt{\frac{-c}{b}}\ \arcsin\left(\frac{s}{\sqrt{p}}\right) + r, \quad r \in \mathrm{I\!R}, \quad p \in \mathrm{I\!R}^*_+.$$

It follows

$$\phi(z) = \pm \frac{a}{c}\sqrt{\frac{-c}{b}}\ \arcsin\left(\frac{1}{e^{\frac{c}{a}z}\sqrt{p}}\right) + r, \quad r \in \mathrm{I\!R}, \quad p \in \mathrm{I\!R}^*_+,$$

that is

$$\phi(z) = \pm \frac{a}{c}\sqrt{\frac{-c}{b}}\ \arcsin\left(Ke^{-\dfrac{c}{a}z}\right) + r, \quad r \in \mathrm{I\!R}, \quad K \in \mathrm{I\!R}^*_+. \tag{19}$$

Therefore, we have found three families of solutions of the multitime Rayleigh PDE (3):

Theorem *a) If we fix the above elements by the conditions*

$$h^{\alpha\beta}\lambda_\alpha\lambda_\beta = constant, \qquad \frac{C^\gamma\lambda_\gamma}{B^{\alpha\beta\gamma}\lambda_\alpha\lambda_\beta\lambda_\gamma} > 0,$$

then we get two families of soliton solutions

$$u(x,t) = \pm \frac{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}{C^\gamma\lambda_\gamma}\sqrt{\frac{C^\gamma\lambda_\gamma}{B^{\alpha\beta\gamma}\lambda_\alpha\lambda_\beta\lambda_\gamma}}\ ch^{-1}\left(Ke^{-\dfrac{C^\gamma\lambda_\gamma}{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}(x - \lambda_\alpha t^\alpha)}\right) + r, \tag{20}$$

where $r \in \mathbb{R},\ \ K \in \mathbb{R}_+^*$, *and*

$$u(x,t) = \pm\frac{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}{C^\gamma\lambda_\gamma}\sqrt{\frac{C^\gamma\lambda_\gamma}{B^{\alpha\beta\gamma}\lambda_\alpha\lambda_\beta\lambda_\gamma}}\, sh^{-1}\left(Ke^{-\dfrac{C^\gamma\lambda_\gamma}{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}(x - \lambda_\alpha t^\alpha)}\right) + r, \tag{21}$$

where $r \in \mathbb{R},\ \ K \in \mathbb{R}_+^*$;

b) if we take

$$h^{\alpha\beta}\lambda_\alpha\lambda_\beta = constant, \qquad \frac{C^\gamma\lambda_\gamma}{B^{\alpha\beta\gamma}\lambda_\alpha\lambda_\beta\lambda_\gamma} < 0,$$

then we have another family of soliton solutions,

$$\begin{aligned} u(x,t) = \pm\, &\frac{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}{C^\gamma\lambda_\gamma}\sqrt{\frac{-C^\gamma\lambda_\gamma}{B^{\alpha\beta\gamma}\lambda_\alpha\lambda_\beta\lambda_\gamma}} \\ &\times \arcsin\left(Ke^{-\dfrac{C^\gamma\lambda_\gamma}{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}(x - \lambda_\alpha t^\alpha)}\right) + r, \end{aligned} \tag{22}$$

where $r \in \mathbb{R},\ \ K \in \mathbb{R}_+^*$.

5.3 Mac-Laurin series soliton of multitime Rayleigh PDE

In a previous section, we obtained the equation (7),

$$a(z)\phi'' - b(z)(\phi')^3 + c(z)\phi' = 0,$$

a second order ODE in ϕ. One assume that it has a solution which is analytic on an interval around $z = 0$ and we search a Mac-Laurin series solution. Then we express ϕ as a power series in the form

$$\phi(z) = \sum_{n=0}^{\infty} \alpha_n\, z^n \tag{23}$$

and we try to determine what the α_n's need to be. The resulting power series need to converge on an interval around origin.

We compute $\phi'(z)$ and $\phi'^3(z)$and $\phi''(z)$:

$$\phi'(z) = \sum_{n=1}^{\infty} n\alpha_n\, z^{n-1} = \sum_{n=0}^{\infty} (n+1)\alpha_{n+1}\, z^n = \sum_{n=0}^{\infty} \beta_n\, z^n,$$

$$(\phi')^2(z) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{n} \beta_k \beta_{n-k} \right) z^n =$$

$$= \sum_{n=0}^{\infty} \left(\sum_{k=0}^{n} \alpha_{k+1}\, \alpha_{n-k+1}(k+1)(n-k+1) \right) z^n = \sum_{n=0}^{\infty} \gamma_n\, z^n,$$

$$(\phi')^3(z) = \sum_{n=0}^{\infty} \left(\sum_{i=0}^{n} \gamma_i\, \beta_{n-i} \right) z^n =$$

$$= \sum_{n=0}^{\infty} \left(\sum_{i=0}^{n} \sum_{k=0}^{i} \alpha_{k+1}\, \alpha_{i-k+1}\, \alpha_{n-i+1}(k+1)(i-k+1)(n-i+1) \right) z^n,$$

$$\phi''(z) = \sum_{n=1}^{\infty} n(n+1)\alpha_{n+1}\, z^{n-1} = \sum_{n=0}^{\infty} (n+1)(n+2)\alpha_{n+2}\, z^n.$$

Consider the particular case

$$a(z) = mz + a, \;\; b(z) = pz + b, \;\; c(z) = qz + c, \;\;\; m, p, q, a, b, c \in \mathbb{R},$$

that is the coefficients $a(z), b(z), c(z)$ of the equation (7) are affine functions in z. Substituting into (7) gives us

$$(mz + a) \sum_{n=0}^{\infty} (n+2)(n+1)\alpha_{n+2}\, z^n + (qz + c) \sum_{n=0}^{\infty} \alpha_{n+1}(n+1)\, z^n -$$

$$-(pz+b) \sum_{n=0}^{\infty} \left(\sum_{i=0}^{n} \sum_{k=0}^{i} \alpha_{k+1}\, \alpha_{i-k+1}\, \alpha_{n-i+1}(k+1)(i-k+1)(n-i+1) \right) z^n = 0.$$

This identity can be written

$$m \sum_{n=1}^{\infty} n(n+1)\alpha_{n+1}\, z^n + 2a\alpha_2 + a \sum_{n=1}^{\infty} (n+2)(n+1)\alpha_{n+2}\, z^n -$$

$$-p \sum_{n=1}^{\infty} \left(\sum_{i=0}^{n-1} \sum_{k=0}^{i} \alpha_{k+1}\, \alpha_{i-k+1}\, \alpha_{n-i}(k+1)(i-k+1)(n-i) \right) z^n - b\alpha_1^3 -$$

$$-b\sum_{n=1}^{\infty}\left(\sum_{i=0}^{n}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i+1}(k+1)(i-k+1)(n-i+1)\right)z^n+$$

$$+q\sum_{n=1}^{\infty}n\alpha_n\,z^n+c\alpha_1+c\sum_{n=1}^{\infty}\alpha_{n+1}(n+1)\,z^n=0,$$

or, equivalent,

$$(2a\alpha_2+c\alpha_1-b\alpha_1^3)+\sum_{n=1}^{\infty}[mn(n+1)\alpha_{n+1}+a(n+2)(n+1)\alpha_{n+2}-$$

$$-p\sum_{i=0}^{n-1}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i}(k+1)(i-k+1)(n-i)-$$

$$-b\sum_{i=0}^{n}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i+1}(k+1)(i-k+1)(n-i+1)+$$

$$+qn\alpha_n+c(n+1)\alpha_{n+1}]z^n=0.$$

By identifying the coefficients of the powers of z with 0, we find the condition

$$2a\alpha_2+c\alpha_1-b\alpha_1^3=0$$

and the recurrence

$$mn(n+1)\alpha_{n+1}+a(n+2)(n+1)\alpha_{n+2}-$$

$$-p\sum_{i=0}^{n-1}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i}(k+1)(i-k+1)(n-i)-$$

$$-b\sum_{i=0}^{n}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i+1}(k+1)(i-k+1)(n-i+1)+$$

$$+qn\alpha_n+c(n+1)\alpha_{n+1}=0,\quad n\geq 1. \tag{24}$$

By the initial conditions $\phi(0)=\alpha_0$, $\phi'(0)=\alpha_1$ and $2a\alpha_2+c\alpha_1-b\alpha_1^3=0$, this recurrence gives us all the coefficients of the power series (23), but the difficult part is just solving the recurrence for the unknown α_n.

Theorem *In the foregoing hypotheses, the multitime series soliton solution of the multitime Rayleigh PDE is*

$$u(x,t)=\sum_{n=0}^{\infty}\alpha_n(x-\lambda_\alpha t^\alpha)^n, \tag{25}$$

with α_0, α_1 *fixed,* $2a\alpha_2 + c\alpha_1 - b\alpha_1^3 = 0$ *and* $\alpha_n, \quad n \geq 2$ *given by the recurrence (24)*

$$mn(n+1)\alpha_{n+1} + a(n+2)(n+1)\alpha_{n+2} -$$
$$-p\sum_{i=0}^{n-1}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i}(k+1)(i-k+1)(n-i)-$$
$$-b\sum_{i=0}^{n}\sum_{k=0}^{i}\alpha_{k+1}\,\alpha_{i-k+1}\,\alpha_{n-i+1}(k+1)(i-k+1)(n-i+1)+$$
$$+qn\alpha_n + c(n+1)\alpha_{n+1} = 0, \quad n \geq 1.$$

6 Families of multitime Rayleigh solitons of Van der Pol type

6.1 Case of coefficients depending on z

As we have seen in a previous section, if we can choose the constant vector λ_α by some conditions which relate the elements h, C, D, λ, then we get the equation

$$a(z)\phi''(z) - d(z)\phi^2(z)\phi'(z) + c(z)\phi'(z) = 0.$$

We can write this equation in the form

$$\phi^2(z)\phi'(z) = \frac{a(z)}{d(z)}\phi''(z) + \frac{c(z)}{d(z)}\phi'(z).$$

In order to find some solutions of this equation, we make a particular choice for λ_α, by a new condition: the coefficients $a(z) \overset{\text{not}}{=} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1$, $d(z) \overset{\text{not}}{=} D^\gamma(t,x,\eta,\xi)\lambda_\gamma$ and $c(z) \overset{\text{not}}{=} C^\gamma(t,x,\eta,\xi)\lambda_\gamma$ will be related by the equality

$$\left(\frac{a(z)}{d(z)}\right)' = \frac{c(z)}{d(z)},$$

or equivalent

$$a'(z)d(z) - a(z)d'(z) = d(z)c(z). \tag{26}$$

With such a selection of λ_α, the equation becomes

$$\phi^2(z)\phi'(z) = \left(\frac{a(z)}{d(z)}\phi'(z)\right)',$$

that is

$$\frac{\phi^3(z)}{3} = \frac{a(z)}{d(z)}\phi'(z) + k, \;\; k \in \mathbb{R}.$$

This Bernoulli ODE has separable variables. By integration, we get

$$\int \frac{3d\phi}{\phi^3 - 3k} = \int \frac{d(z)}{a(z)}\, dz, \;\; k \in \mathbb{R}. \tag{27}$$

For simplicity, we can take $k = 0$ and we find a particular family of solutions, defined by the relation

$$\phi^2(z) = -\frac{2}{3} \int \frac{d(z)}{a(z)}\, dz.$$

If we keep k variable and non-zero, then making the substitution $3k$ by k_1^3, the equality (27) becomes

$$\int \frac{3d\phi}{\phi^3 - k_1^3} = \int \frac{d(z)}{a(z)}\, dz, \;\; k_1 \in \mathbb{R}^*.$$

The integral from the left hand is

$$\int \frac{3d\phi}{(\phi - k_1)(\phi^2 + \phi k_1 + k_1^2)} = \frac{1}{k_1^2} \int \frac{d\phi}{\phi - k_1} - \frac{1}{k_1^2} \int \frac{\phi + 2k_1}{\phi^2 + \phi k_1 + k_1^2}\, d\phi =$$

$$= \frac{1}{k_1^2} \ln|\phi - k_1| - \frac{1}{2k_1^2} \int \frac{2\phi + k_1}{\phi^2 + \phi k_1 + k_1^2}\, d\phi - \frac{3}{2k_1} \int \frac{d\phi}{\left(\phi + \frac{k_1}{2}\right)^2 + \frac{3k_1^2}{4}} =$$

$$= \frac{1}{k_1^2} \ln|\phi - k_1| - \frac{1}{2k_1^2} \ln(\phi^2 + \phi k_1 + k_1^2) - \frac{\sqrt{3}}{k_1^2} \operatorname{arctg}\left(\frac{2\phi + k_1}{k_1\sqrt{3}}\right) + c_1, \;\; c_1 \in \mathbb{R}.$$

Therefore, the general solution of the equation (27) is expressed implicitly by the equality

$$\frac{1}{k_1^2} \ln \frac{|\phi - k_1|}{\sqrt{\phi^2 + \phi k_1 + k_1^2}} - \frac{\sqrt{3}}{k_1^2} \operatorname{arctg}\left(\frac{2\phi + k_1}{k_1\sqrt{3}}\right) = \int \frac{d(z)}{a(z)}\, dz, \;\; k_1 \in \mathbb{R}^*.$$

Summarizing, we can formulate the next result:

Theorem *If we can take the constant vector* λ_α *so as the elements* $h^{\alpha\beta}(t, x, \eta, \xi)$, $B^\gamma(t, x, \eta, \xi)$, $C^\gamma(t, x, \eta, \xi)$ *and* λ_α, $\alpha, \beta, \gamma = 1, ..., m$, *to be* *related by the conditions*

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = a(x-\lambda_\alpha t^\alpha) \neq 0, \\ B^\gamma(t,x,\eta,\xi)\lambda_\gamma = b(x-\lambda_\alpha t^\alpha), \\ C^\gamma(t,x,\eta,\xi)\lambda_\gamma = c(x-\lambda_\alpha t^\alpha), \\ a'(z)d(z) - a(z)b'(z) = d(z)c(z), \end{cases}$$

then the function

$$u(x,t) = \phi(x-\lambda_\alpha t^\alpha)$$

represents a multitime soliton-solution for the multitime PDE Van der Pol (3'), for every ϕ *defined implicitly by one of the equalities*

$$\phi^2(z) = -\frac{2}{3}\int \frac{d(z)}{a(z)}\,dz \tag{28}$$

or

$$\frac{1}{k_1^2}\ln\frac{|\phi(z)-k_1|}{\sqrt{\phi^2(z)+\phi(z)k_1+k_1^2}} - \frac{\sqrt{3}}{k_1^2}arctg\left(\frac{2\phi(z)+k_1}{k_1\sqrt{3}}\right) = \int \frac{d(z)}{a(z)}\,dz, \quad k_1 \in \mathbb{R}^*. \tag{29}$$

6.2 Case of constant coefficients

If we fix the elements h, C, D, λ by the conditions

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = \text{constant} = a \neq 0, \\ D^\gamma(t,x,\eta,\xi)\lambda_\gamma = \text{constant} = d, \\ C^\gamma(t,x,\eta,\xi)\lambda_\gamma = \text{constant} = c, \end{cases}$$

then the ODE (7') takes the form

$$a\,\phi'' - d\,\phi^2\phi' + c\,\phi' = 0. \tag{30}$$

Integrating, we find a first order ODE, with separable variables

$$a\,\phi' - \frac{d}{3}\phi^3 + c\phi = k, \;\; k \in \mathbb{R}.$$

In order to find some solutions of this equation, we take $k = 0$. By this choice, the equation becomes a *Bernoulli* ODE. We make a change of the unknown function,

$$\psi = \phi^{-2},$$

and the Bernoulli ODE becomes a linear ODE,

$$\psi' - \frac{2c}{a}\psi = -\frac{2d}{3a},$$

with the solutions

$$\psi(z) = e^{\int \frac{2c}{a} dz} \left[K - \int \frac{2d}{3a} e^{-\int \frac{2c}{a} dz} dz \right], \quad K \in \mathbb{R},$$

that is

$$\psi(z) = K e^{\frac{2c}{a} z} + \frac{d}{3c}, \quad K \in \mathbb{R}. \tag{31}$$

Since $\phi = \psi^{-\frac{1}{2}}$, we obtain

$$\phi(z) = \frac{1}{\sqrt{K e^{\frac{2c}{a} z} + \frac{d}{3c}}}, \quad K \in \mathbb{R}. \tag{32}$$

Thus we obtain a family of multitime soliton-solutions of the multitime Van der Pol PDE (3') and we can formulate the next theorem:

Theorem *If we take* λ_α *so as to fix* h, D, C, λ *by the conditions*

$$\begin{cases} h^{\alpha\beta}(t,x,\eta,\xi)\lambda_\alpha\lambda_\beta - 1 = constant = a \neq 0, \\ D^\gamma(t,x,\eta,\xi)\lambda_\gamma = constant = d, \\ C^\gamma(t,x,\eta,\xi)\lambda_\gamma = constant = c, \end{cases}$$

then we get a family of multitime soliton-solutions of the multitime Rayleigh wave equation of Van der Pol type $(3')$,

$$u(x,t) = \frac{1}{\sqrt{K e^{\frac{2C^\gamma\lambda_\gamma}{h^{\alpha\beta}\lambda_\alpha\lambda_\beta - 1}(x - \lambda_\alpha t^\alpha)} + \frac{D^\gamma\lambda_\gamma}{3(C^\gamma\lambda_\gamma)}}}, \quad K \in \mathbb{R}. \tag{33}$$

Acknowledgements. Partially supported by University Politehnica of Bucharest, and by Academy of Romanian Scientists, Bucharest, Romania.

References

[1] J. W. S. Rayleigh, *On waves propagating along the plane surface of an elastic solid*, Proc. London Math. Soc., 17 (1887), 4-11.

[2] Darmawijoyo, W. T. van Horssen, P. Clement - *On a Rayleigh wave equation with boundary damping*, Nonlinear Dynamics, 33, 4 (2003), 399-429.

[3] W. S. Hall - *The Rayleigh and Van der Pol wave equations, some generalizations*, Lecture Notes in Mathematics, 703 (1979), 103-138, Springer-Verlag, Berlin.

[4] P. G. Malischewsky, Comment to "A new formula for the velocity of Rayleigh waves" by D.Nkemzi, Wave Motion, 26 (1997), 199-205]; Wave Motion, 31 (2000), 93-96.

[5] L. Matei, C. Udrişte, *Multitime sine-Gordon solitons via geometric characteristics*, in Balkan Journal of Geometry and Its Applications, 16, 2 (2011), 81-89.

[6] Houari Mechkour, The exact expressions for the roots of Rayleigh wave equation, BSG Proceedings, 8 (2003), 96 - 104, Geometry Balkan Press, Bucharest.

[7] D. Nkemzi, *A new formula for the velocity of Rayleigh waves*, Wave Motion, 26 (1997), 199-205.

[8] M. Rahman, J. R. Barber, *Exact expressions for the roots of the secular equation for Rayleigh waves*, ASME, J. Appl. Mech., 62 (1995) 250-252.

[9] D. Royer, *A study of the secular equation for Rayleigh waves using the root locus method*, Ultrasonics, 39 (2001), 223-225.

[10] C. Udrişte, M. Ferrara, D. Opriş, *Economic Geometric Dynamics*, Geometry Balkan Press, Bucharest, 2004.

[11] C. Udrişte, *Simplified multitime maximum principle*, Balkan Journal of Geometry and Its Applications, 14, 1 (2009), 102-119.

[12] C. Udrişte, *Nonholonomic approach of multitime maximum principle*, Balkan Journal of Geometry and Its Applications, 14, 2 (2009), 111-126.

[13] C. Udrişte, *Multitime controllability, observability and bang-bang principle*, Journal of Optimization Theory and Applications, 139, 1 (2008), 141-157.

[14] C. Udrişte, I. Ţevy, *Multitime Linear-Quadratic Regulator Problem Based on Curvilinear Integral*, Balkan Journal of Geometry and Its Applications, 14, 2 (2009), 127-137.

[15] C. Udrişte, I. Ţevy, *Multitime Dynamic Programming for Curvilinear Integral Actions*, Journal of Optimization Theory and Applications, 146 (2010), 189-207.

[16] C. Udrişte, *Equivalence of multitime optimal control problems*, Balkan Journal of Geometry and Its Applications, 15, 1 (2010), 155-162.

[17] C. Udrişte, L. Matei, I. Duca, *Multitime Hamilton-Jacobi Theory*, Proceedings of 8-th WSEAS International Conference on Applied Computational Science (ACACOS-09), 509-513, Hangzhou, China, May 20-22, 2009; Proceedings of the 9-th WSEAS International Conference on Systems Theory and Scientific Computation (ISTASC-09), 129-133, Moscow, Russia, August 20-22, 2009.